

PhD Project Proposal 7

Singular Integral Operators in Potential Theory: at the Analysis/Numerical Analysis Interface

Supervisors: [Prof Simon Chandler-Wilde](#), [Prof Jani Virtanen](#), and [Dr Raffael Hagger](#)

In this project we will attack a long-standing conjecture in potential theory of (Kenig, 1994) regarding the so-called *double-layer potential operator* K on a closed curve $\Gamma \subset \mathbb{R}^2$. The conjecture is that K has essential spectral radius $< \frac{1}{2}$ whenever Γ is the boundary of a bounded Lipschitz domain.

To explain these terms, let $G(x, y) := \log(|x - y|) / (2\pi)$, for $x, y \in \mathbb{R}^2$, the so-called *fundamental solution of Laplace's equation*, and suppose $D \subset \mathbb{R}^2$ is a bounded Lipschitz domain with boundary Γ , meaning that, at each $x \in \Gamma$, Γ is locally the graph of some Lipschitz continuous function in some rotated coordinate system. Then the double-layer potential operator K on $L^2(\Gamma, ds)$ (the set of functions on Γ that are square integrable with respect to arc-length measure ds on Γ) is defined by

$$K\phi(x) = \int_{\Gamma} \frac{\partial G(x, y)}{\partial n(y)} \phi(y) ds(y), \quad x \in \Gamma,$$

for every $\phi \in L^2(\Gamma, ds)$, where the derivative is in the direction of the unit normal $n(y)$ at y . The spectrum of an operator K is the set of values $\lambda \in \mathbb{C}$ for which $\lambda I - K$ is not invertible, where I is the identity operator. The essential spectrum, $\text{spec}_{\text{ess}} K$, is arguably the most important part of the spectrum for a bounded operator on a Hilbert space, and is the set of $\lambda \in \mathbb{C}$ for which $\lambda I - K$ is not

even approximately invertible in a sense that can be made precise.

Kenig's conjecture has been shown to hold when Γ is the boundary of a convex domain, and when Γ is a polygon. In this project we will explore whether the conjecture holds for much more complicated boundaries, for example the self-similar boundary Γ shown in Figure 1.




Figure 1. An example of a complicated Lipschitz boundary.

To do this we will use a combination of methods which will depend on the background and interests of the candidate but will include analysis/functional analysis methods and rigorously justified numerical analysis and computation of $\text{spec}_{\text{ess}} K$ in MATLAB or Python.

In carrying out this work the candidate will work with and learn from the supervisors and will also have opportunities to interact with and visit other leading experts in this area, for example [Euan Spence](#) (Bath) and [Karl-Mikael Perfekt](#) (Trondheim), and will present their progress at international conferences. For more information about the topic see the [recent preprint](#) (Chandler-Wilde & Spence, 2021).

Chandler-Wilde, S. N., & Spence, E. A. (2021). Coercivity, essential norms, and the Galerkin method for second-kind integral equations on polyhedral and Lipschitz domains. Retrieved from <https://arxiv.org/abs/2105.11383>

Kenig, C. E. (1994). *Harmonic analysis techniques for second order elliptic boundary value problems*. American Mathematical Society.